



11. Current research:

ONS Foundation-funded research

<http://www.ons.org/research/funding/Projects/index.shtml>

“Symptom Clusters in Children/Teens on Cisplatin or Ifosfamide,” Marilyn Hockenberry, PhD, RN, Baylor College of Medicine

Cancer Research-UK <http://www.ncri.org.uk/home/index.cfm>

Tamoxifen Chemoprevention Trial

National Institutes of Health – funded research <http://www.nih.gov/>

Development of Poly-L-Glutamic Acid Paclitaxel Conjugate (National Cancer Institute [NCI])

Effects of Acupuncture on Pain, Nausea, Quality of Life (NCI)

Impact of Hypnosis for Breast Surgery Side Effects (NCI)

Presurgery Hypnosis--Benefits Analysis in Breast Cancer (NCI)

Treating Chemotherapy Induced Nausea with Acupressure (NCI)

Symptom Clusters in Cancer Patients Undergoing Treatment (National Institute of Nursing Research)

Psychoimmune Outcomes: Intervention in Breast Cancer

TNF Blockade in Pancreatic Cancer Patients (NCI)

Ginger Control of Chemotherapy Induced Nausea and Emesis (National Center for Complementary and Alternative Medicine)

Department of Defense: Congressionally Directed Medical Research Programs

<http://cdmmp.army.mil/default.htm>

A Controlled Study Using Acupuncture as an Adjuvant to Treat Chemotherapy-Induced Nausea and Vomiting Acustimulation for the Control of Chemotherapy-Induced Nausea in Breast Cancer

http://www.cancer.org/docroot/RES/RES_0.asp

American Cancer Society and National Comprehensive Cancer Network. (2001). Preventive treatment, breakthrough treatment, subsequent cycles, delayed nausea and vomiting, and anticipatory nausea and vomiting. In *Nausea and vomiting treatment guidelines for patients with cancer* (pp. 5–6). Retrieved July 23, 2004, from [Nausea and Vomiting Treatment Guidelines for Patients with Cancer](#)

Alves-Guerreiro, J., Lowe-Strong, A.S., Walsh, D.M., Lopes, B.C., Costa, R., Rosado, R., et al. (2003). Development of a Portuguese Version of the Morrow Assessment of Nausea and Emesis (MANE) Questionnaire; Moving physical therapy forward [Abstract]. *14th International World Confederation for Physical Therapy Congress*

- Arakawa, S. (1995). Effectiveness of progressive muscle relaxation in reducing nausea, vomiting, and anxiety induced by chemotherapy in Japanese patients. *Dissertation Abstracts International: Section B: the Sciences & Engineering*, 56(5).
- Arakawa, S. (1997). Relaxation to reduce nausea, vomiting, and anxiety induced by chemotherapy in Japanese patients. *Cancer Nursing*, 20, 342–349.
- ASHP therapeutic guidelines on the pharmacologic management of nausea and vomiting in adult and pediatric patients receiving chemotherapy or radiation therapy or undergoing surgery. (1999). *American Journal of Health System Pharmacists*, 56, 729–764
- Bonneterre, J., Schraub, S., Lecomte, S., & Mercier, M. (1996). Quality of life as an outcome in breast cancer. *Pharmacoeconomics*, 9 Suppl 2, 23–29.
- Crucitt, M.A., Hyman, W., Grote, T., Tester, W., Madajewicz, S., Yee, S., et al. (1996). Efficacy and tolerability of oral ondansetron versus prochlorperazine in the prevention of emesis associated with cyclophosphamide-based chemotherapy and maintenance of health related quality of life. *Clinical Therapeutics*, 18, 778–788.
- del Giglio, A., Soares, H.P., Caparroz, C., & Castro P.C. (2000). Granisetron is equivalent to ondansetron for prophylaxis of chemotherapy-induced nausea and vomiting: results of a meta-analysis of randomized controlled trials. *Cancer*, 89, 2301–2308.
- Devine, E.C. & Westlake, S.K. (1995). The effects of psychoeducational care provided to adults with cancer: meta-analysis of 116 studies. *Oncology Nursing Forum*, 22, 1369–1381. [Oncology Nursing Forum](#)
- Dibble, S.L., Chapman, J., Mack, K.A., Shih, A. (2000). Acupressure for nausea: Results of a pilot study. *Oncology Nursing Forum*, 27, 41–47. [Oncology Nursing Forum](#)
- Dodd, M.J., Onishi, K., Dibble, S.L., & Larson, P.J. (1996). Differences in nausea, vomiting, and retching between younger and older outpatients receiving cancer chemotherapy. *Cancer Nursing*, 19, 155–161.
- Farley, P.A., Dempsey, C.L., Shillington, A.A., Kulis-Robitaille, C., Colgan, K., & Bernstein, G. (1997). Patients' self-reported functional status after granisetron or ondansetron therapy to prevent chemotherapy-induced nausea and vomiting at six cancer centers. *American Journal of Health-System Pharmacy*, 54, 2478–2482.
- Fazeny-Dorner, B., Veitl, M., Wenzel, C., Brodowicz, T., Zielinski, C., Muhm M. et al. (2002). Alterations in intestinal permeability following the intensified polydrug-chemotherapy IFADIC (ifosfamide, Adriamycin, dacarbazine). *Cancer Chemotherapy and Pharmacology*, 49, 294–298.
- Fu, M.R., Rhodes, V., & Xu, B. (2002). The Chinese translation of the Index of Nausea, Vomiting, and Retching. *Cancer Nursing*, 25, 134–140.
- Gralla, R.J., Osoba, D., Kris, M.G., Kirkbride, P., Hesketh, P.J., Chinnery, L.W., et al. (1999). Recommendations for the use of antiemetics: Evidence-based, clinical practice guidelines. American Society of Clinical Oncology. *Journal of Clinical Oncology*, 17, 2971–2994

- Gilbert, C.J., Ohly, K.V., Rosner, G., & Peters, W.P. (1995). Randomized, double-blind comparison of a prochlorperazine-based versus a metoclopramide-based antiemetic regimen in patients undergoing autologous bone marrow transplantation. *Cancer*, *76*, 2330–2337.
- Hickok, J.T., Roscoe, J.A., Morrow, G.R., Stern, R.M., Yang, B., Flynn, P.J., et al. (1999). Use of 5-HT₃ receptor antagonists to prevent nausea and emesis caused by chemotherapy for patients with breast carcinoma in community practice settings. *Cancer*, *86*, 64–71.
- Ioannidis, J.P.A., Hesketh, P.J., & Lau, J. (2000). Contribution of dexamethasone to control of chemotherapy-induced nausea and vomiting: A meta-analysis of randomized evidence. *Journal of Clinical Oncology*, *18*, 3409–3422.
- Lebeau, B., Depierre, A., Giovannini, M., Riviere, A., Kaluzinski, L., Votan B., et al. (1997). The efficacy of a combination of ondansetron, methylprednisolone and metopimazine in patients previously uncontrolled with a dual antiemetic treatment in cisplatin-based chemotherapy. The French Ondansetron Study Group. *Annals of Oncology*, *8*, 887–892.
- Lindley, C.M., Hirsch, J.D., O'Neill, C.V., Transau, M.C., Gilbert, C.S., & Osterhaus, J.T. (1992). Quality of life consequences of chemotherapy-induced emesis. *Quality of Life Research*, *1*, 331–340.
- Lo, L.H. & Hayman, L.L. (1999). Parents associated with children in measuring acute and delayed nausea and vomiting. *Nursing and Health Sciences*, *1*, 155–161.
- Mantovani, G., Astara, G., Lampis, B., Bianchi, A., Curreli, L., Orru, W., et al. (1996). Evaluation by multidimensional instruments of health-related quality of life of elderly cancer patients undergoing three different "psychosocial" treatment approaches. A randomized clinical trial. *Supportive Care in Cancer*, *4*, 129–140.
- Martin, A.R., Carides, A.D., Pearson, J.D., Horgan, K., Elmer, M., Schmidt, C. et al. (2003). Functional relevance of antiemetic control. Experience using the FLIE questionnaire in a randomised study of the NK-1 antagonist aprepitant. *European Journal of Cancer*, *39*, 1395-1401.
- Martin, C.G., Rubenstein, E.B., Elting, L.S., Kim, Y.J., & Osoba, D. (2003). Measuring chemotherapy-induced nausea and emesis. *Cancer*, *98*, 645–655.
- Ming, J.L., Kuo, B.I., Lin J.G., & Lin, L.C. (2002). The efficacy of acupuncture to prevent nausea and vomiting in post-operative patients. *Journal of Advanced Nursing*, *39*, 343–351.
- Molassiotis, A. (2000). A pilot study of the use of progressive muscle relaxation training in the management of post-chemotherapy nausea and vomiting. *European Journal of Cancer Care*, *9*, 230–234.
- Molassiotis, A., Yung, H.P., Yam, B.M., Chan, F.Y., & Mok, T.S. (2002). The effectiveness of progressive muscle relaxation training in managing chemotherapy-induced nausea and vomiting in Chinese breast cancer patients: A randomised controlled trial. *Supportive Care in Cancer*, *10*, 237–246

- Morrow, G. (1984). Assessment of nausea and vomiting: Past problems, current issues and suggestions for future research. *Cancer*, 53, 2267–2278.
- Morrow, G.R. (1992). Methodology and assessment in clinical anti-emetic research: A meta analysis of outcome parameters. *British Journal of Cancer*, 19(Suppl.), S38–41.
- Morrow, G.R., Roscoe, J.A., Hickok, J.T., Stern, R.M., Pierce, H.I., King, D.B., et al. (1998). Initial control of chemotherapy-induced nausea and vomiting in patient quality of life. *Oncology*, 12(3, Suppl. 4), 32–37.
- Mullin, S.M., Fletcher, D.M. & Tyler, L.S. (1998). Mail-in questionnaire for monitoring nausea and vomiting in oncology outpatients. *American Journal of Health-System Pharmacy*, 55, 1903–1906.
- Pan, C.X., Morrison, R.S., Ness, J., Fugh-Berman, A., & Leipzig, R.M. (2000). Complementary and alternative medicine in the management of pain, dyspnea, and nausea and vomiting near the end of life: A systematic review. *Journal of Pain and Symptom Management*, 20, 374–387.
- Rhodes, V.A., & McDaniel R.W. (1999). The Index of Nausea, Vomiting, and Retching: A new format of the Index of Nausea and Vomiting. *Oncology Nursing Forum*, 26, 889–894. [Oncology Nursing Forum](#)
- Roscoe, J.A., Morrow, G.R., Hickok, J.T., & Stern, R.M. (2000). Nausea and vomiting remain a significant clinical problem: trends over time in controlling chemotherapy-induced nausea and vomiting in 1413 patients treated in community clinical practices. *Journal of Pain & Symptom Management*, 20, 113–121.
- Satou, A., Yamazaki, T., Nukariya, N., Nakamachi, M., Shimada, K., Matsukawa, M., et al. (2002). Development of a Japanese version of the FLIE. *Gan to Kagaku Ryoho [Japanese Journal of Cancer and Chemotherapy]*, 29, 281–291.
- Sloan, J.A., Goldberg, R.M., Sargent, D.J., Vargas-Chanes, D., Nair, S. Cha, S.S., et al. (2002). Women experience greater toxicity with fluorouracil-based chemotherapy for colorectal cancer. *Journal of Clinical Oncology*, 20, 1491–1498.
- Smith, M.C., Holcombe, J.K., & Stullenbarger, E. (1994). A meta-analysis of intervention effectiveness for symptom management in oncology nursing research. *Oncology Nursing Forum*, 21, 1201–1210.
- Sykes, A.J., Kiltie, A.E., & Stewart, A.L. (1997). Ondansetron versus a chlorpromazine And dexamethasone combination for the prevention of nausea and vomiting: a prospective, randomised study to assess efficacy, cost effectiveness and quality of life following single-fraction radiotherapy. *Supportive Care in Cancer*, 5, 500–503.
- Tramer, M. R, Carroll, D., Campbell, F.A., Reynolds, D.J, Moore, R.A., & McQuay, H.J. (2003). Cannabinoids for control of chemotherapy induced nausea and vomiting: Quantitative systematic review. *BMJ*, 323(7303), 16–21.
- World Health Organization. (1979). WHO Handbook for Reporting Results of Cancer Treatment. Geneva: Author. Retrieved July 23, 2004, from [WHO Handbook for Reporting Results of Cancer Treatment](#)